





INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(51) International Patent Classification⁴ : A61K 37/02, C07K 5/06, 5/08 C07K 5/10, C07D 405/00 C07D 207/00	A1	(11) International Publication Number: WO 89/ 03223 (43) International Publication Date: 20 April 1989 (20.04.89)
(21) International Application Number: PCT/US88/03506 (22) International Filing Date: 6 October 1988 (06.10.88) (31) Priority Application Number: 105,768 (32) Priority Date: 6 October 1987 (06.10.87) (33) Priority Country: US (71)(72) Applicants and Inventors: BACHOVCHIN, William, W. [US/US]; 71 Warwick Road, Melrose, MA 02176 (US). PLAUT, Andrew, G. [US/US]; 22 Peacock Farm Road, Lexington, MA 02173 (US). KETTER, Charles, A. [US/US]; 2411 Chatham Drive, Wilmington, DE 19803 (US). (74) Agent: CLARK, Paul, T.; Fish & Richardson, One Financial Center, Suite 2500, Boston, MA 02111-2658 (US).		(81) Designated States: AT (European patent), AU, BE (European patent), CH (European patent), DE (European patent), DK, FI, FR (European patent), GB (European patent), IT (European patent), JP, LU (European patent), NL (European patent), NO, SE (European patent). Published <i>With international search report.</i> <i>Before the expiration of the time limit for amending the claims and to be republished in the event of the receipt of amendments.</i>


Abandoned

(54) Title: PROTEASE INHIBITORS


Prolyl Boronate




II



III




Prolyl phosphonate





IV



V



VI



VII

group of formula (I) where G is either H or an alkyl group containing 1 to about 20 carbon atoms and optional heteroatoms which can be N, S, or O; or a phosphonate group of formula (IV) where J is O-alkyl, N-alkyl, or alkyl, each comprising about 1-20 carbon atoms and, optionally, heteroatoms which can be N, S, or O; T being able to form a complex with the catalytic site of an enzyme. X is a group having at least one amino acid. Y is (V) or (VI) or (VII) and each R¹, R², R³, R⁴, R⁵, R⁶, R⁷, R⁸, R⁹, R¹⁰, R¹¹, R¹², R¹³, R¹⁴, R¹⁵, R¹⁶, R¹⁷, R¹⁸, R¹⁹, R²⁰, R²¹, R²², R²³, R²⁴, R²⁵, R²⁶, R²⁷, R²⁸, R²⁹, R³⁰, R³¹, R³², R³³, R³⁴, R³⁵, R³⁶, R³⁷, R³⁸, R³⁹, R⁴⁰, R⁴¹, R⁴², R⁴³, R⁴⁴, R⁴⁵, R⁴⁶, R⁴⁷, R⁴⁸, R⁴⁹, R⁵⁰, R⁵¹, R⁵², R⁵³, R⁵⁴, R⁵⁵, R⁵⁶, R⁵⁷, R⁵⁸, R⁵⁹, R⁶⁰, R⁶¹, R⁶², R⁶³, R⁶⁴, R⁶⁵, R⁶⁶, R⁶⁷, R⁶⁸, R⁶⁹, R⁷⁰, R⁷¹, R⁷², R⁷³, R⁷⁴, R⁷⁵, R⁷⁶, R⁷⁷, R⁷⁸, R⁷⁹, R⁸⁰, R⁸¹, R⁸², R⁸³, R⁸⁴, R⁸⁵, R⁸⁶, R⁸⁷, R⁸⁸, R⁸⁹, R⁹⁰, R⁹¹, R⁹², R⁹³, R⁹⁴, R⁹⁵, R⁹⁶, R⁹⁷, R⁹⁸, R⁹⁹, R¹⁰⁰, R¹⁰¹, R¹⁰², R¹⁰³, R¹⁰⁴, R¹⁰⁵, R¹⁰⁶, R¹⁰⁷, R¹⁰⁸, R¹⁰⁹, R¹¹⁰, R¹¹¹, R¹¹², R¹¹³, R¹¹⁴, R¹¹⁵, R¹¹⁶, R¹¹⁷, R¹¹⁸, R¹¹⁹, R¹²⁰, R¹²¹, R¹²², R¹²³, R¹²⁴, R¹²⁵, R¹²⁶, R¹²⁷, R¹²⁸, R¹²⁹, R¹³⁰, R¹³¹, R¹³², R¹³³, R¹³⁴, R¹³⁵, R¹³⁶, R¹³⁷, R¹³⁸, R¹³⁹, R¹⁴⁰, R¹⁴¹, R¹⁴², R¹⁴³, R¹⁴⁴, R¹⁴⁵, R¹⁴⁶, R¹⁴⁷, R¹⁴⁸, R¹⁴⁹, R¹⁵⁰, R¹⁵¹, R¹⁵², R¹⁵³, R¹⁵⁴, R¹⁵⁵, R¹⁵⁶, R¹⁵⁷, R¹⁵⁸, R¹⁵⁹, R¹⁶⁰, R¹⁶¹, R¹⁶², R¹⁶³, R¹⁶⁴, R¹⁶⁵, R¹⁶⁶, R¹⁶⁷, R¹⁶⁸, R¹⁶⁹, R¹⁷⁰, R¹⁷¹, R¹⁷², R¹⁷³, R¹⁷⁴, R¹⁷⁵, R¹⁷⁶, R¹⁷⁷, R¹⁷⁸, R¹⁷⁹, R¹⁸⁰, R¹⁸¹, R¹⁸², R¹⁸³, R¹⁸⁴, R¹⁸⁵, R¹⁸⁶, R¹⁸⁷, R¹⁸⁸, R¹⁸⁹, R¹⁹⁰, R¹⁹¹, R¹⁹², R¹⁹³, R¹⁹⁴, R¹⁹⁵, R¹⁹⁶, R¹⁹⁷, R¹⁹⁸, R¹⁹⁹, R²⁰⁰, R²⁰¹, R²⁰², R²⁰³, R²⁰⁴, R²⁰⁵, R²⁰⁶, R²⁰⁷, R²⁰⁸, R²⁰⁹, R²¹⁰, R²¹¹, R²¹², R²¹³, R²¹⁴, R²¹⁵, R²¹⁶, R²¹⁷, R²¹⁸, R²¹⁹, R²²⁰, R²²¹, R²²², R²²³, R²²⁴, R²²⁵, R²²⁶, R²²⁷, R²²⁸, R²²⁹, R²³⁰, R²³¹, R²³², R²³³, R²³⁴, R²³⁵, R²³⁶, R²³⁷, R²³⁸, R²³⁹, R²⁴⁰, R²⁴¹, R²⁴², R²⁴³, R²⁴⁴, R²⁴⁵, R²⁴⁶, R²⁴⁷, R²⁴⁸, R²⁴⁹, R²⁵⁰, R²⁵¹, R²⁵², R²⁵³, R²⁵⁴, R²⁵⁵, R²⁵⁶, R²⁵⁷, R²⁵⁸, R²⁵⁹, R²⁶⁰, R²⁶¹, R²⁶², R²⁶³, R²⁶⁴, R²⁶⁵, R²⁶⁶, R²⁶⁷, R²⁶⁸, R²⁶⁹, R²⁷⁰, R²⁷¹, R²⁷², R²⁷³, R²⁷⁴, R²⁷⁵, R²⁷⁶, R²⁷⁷, R²⁷⁸, R²⁷⁹, R²⁸⁰, R²⁸¹, R²⁸², R²⁸³, R²⁸⁴, R²⁸⁵, R²⁸⁶, R²⁸⁷, R²⁸⁸, R²⁸⁹, R²⁹⁰, R²⁹¹, R²⁹², R²⁹³, R²⁹⁴, R²⁹⁵, R²⁹⁶, R²⁹⁷, R²⁹⁸, R²⁹⁹, R³⁰⁰, R³⁰¹, R³⁰², R³⁰³, R³⁰⁴, R³⁰⁵, R³⁰⁶, R³⁰⁷, R³⁰⁸, R³⁰⁹, R³¹⁰, R³¹¹, R³¹², R³¹³, R³¹⁴, R³¹⁵, R³¹⁶, R³¹⁷, R³¹⁸, R³¹⁹, R³²⁰, R³²¹, R³²², R³²³, R³²⁴, R³²⁵, R³²⁶, R³²⁷, R³²⁸, R³²⁹, R³³⁰, R³³¹, R³³², R³³³, R³³⁴, R³³⁵, R³³⁶, R³³⁷, R³³⁸, R³³⁹, R³⁴⁰, R³⁴¹, R³⁴², R³⁴³, R³⁴⁴, R³⁴⁵, R³⁴⁶, R³⁴⁷, R³⁴⁸, R³⁴⁹, R³⁵⁰, R³⁵¹, R³⁵², R³⁵³, R³⁵⁴, R³⁵⁵, R³⁵⁶, R³⁵⁷, R³⁵⁸, R³⁵⁹, R³⁶⁰, R³⁶¹, R³⁶², R³⁶³, R³⁶⁴, R³⁶⁵, R³⁶⁶, R³⁶⁷, R³⁶⁸, R³⁶⁹, R³⁷⁰, R³⁷¹, R³⁷², R³⁷³, R³⁷⁴, R³⁷⁵, R³⁷⁶, R³⁷⁷, R³⁷⁸, R³⁷⁹, R³⁸⁰, R³⁸¹, R³⁸², R³⁸³, R³⁸⁴, R³⁸⁵, R³⁸⁶, R³⁸⁷, R³⁸⁸, R³⁸⁹, R³⁹⁰, R³⁹¹, R³⁹², R³⁹³, R³⁹⁴, R³⁹⁵, R³⁹⁶, R³⁹⁷, R³⁹⁸, R³⁹⁹, R⁴⁰⁰, R⁴⁰¹, R⁴⁰², R⁴⁰³, R⁴⁰⁴, R⁴⁰⁵, R⁴⁰⁶, R⁴⁰⁷, R⁴⁰⁸, R⁴⁰⁹, R⁴¹⁰, R⁴¹¹, R⁴¹², R⁴¹³, R⁴¹⁴, R⁴¹⁵, R⁴¹⁶, R⁴¹⁷, R⁴¹⁸, R⁴¹⁹, R⁴²⁰, R⁴²¹, R⁴²², R⁴²³, R⁴²⁴, R⁴²⁵, R⁴²⁶, R⁴²⁷, R⁴²⁸, R⁴²⁹, R⁴³⁰, R⁴³¹, R⁴³², R⁴³³, R⁴³⁴, R⁴³⁵, R⁴³⁶, R⁴³⁷, R⁴³⁸, R⁴³⁹, R⁴⁴⁰, R⁴⁴¹, R⁴⁴², R⁴⁴³, R⁴⁴⁴, R⁴⁴⁵, R⁴⁴⁶, R⁴⁴⁷, R⁴⁴⁸, R⁴⁴⁹, R⁴⁵⁰, R⁴⁵¹, R⁴⁵², R⁴⁵³, R⁴⁵⁴, R⁴⁵⁵, R⁴⁵⁶, R⁴⁵⁷, R⁴⁵⁸, R⁴⁵⁹, R⁴⁶⁰, R⁴⁶¹, R⁴⁶², R⁴⁶³, R⁴⁶⁴, R⁴⁶⁵, R⁴⁶⁶, R⁴⁶⁷, R⁴⁶⁸, R⁴⁶⁹, R⁴⁷⁰, R⁴⁷¹, R⁴⁷², R⁴⁷³, R⁴⁷⁴, R⁴⁷⁵, R⁴⁷⁶, R⁴⁷⁷, R⁴⁷⁸, R⁴⁷⁹, R⁴⁸⁰, R⁴⁸¹, R⁴⁸², R⁴⁸³, R⁴⁸⁴, R⁴⁸⁵, R⁴⁸⁶, R⁴⁸⁷, R⁴⁸⁸, R⁴⁸⁹, R⁴⁹⁰, R⁴⁹¹, R⁴⁹², R⁴⁹³, R⁴⁹⁴, R⁴⁹⁵, R⁴⁹⁶, R⁴⁹⁷, R⁴⁹⁸, R⁴⁹⁹, R⁵⁰⁰, R⁵⁰¹, R⁵⁰², R⁵⁰³, R⁵⁰⁴, R⁵⁰⁵, R⁵⁰⁶, R⁵⁰⁷, R⁵⁰⁸, R⁵⁰⁹, R⁵¹⁰, R⁵¹¹, R⁵¹², R⁵¹³, R⁵¹⁴, R⁵¹⁵, R⁵¹⁶, R⁵¹⁷, R⁵¹⁸, R⁵¹⁹, R⁵²⁰, R⁵²¹, R⁵²², R⁵²³, R⁵²⁴, R⁵²⁵, R⁵²⁶, R⁵²⁷, R⁵²⁸, R⁵²⁹, R⁵³⁰, R⁵³¹, R⁵³², R⁵³³, R⁵³⁴, R⁵³⁵, R⁵³⁶, R⁵³⁷, R⁵³⁸, R⁵³⁹, R⁵⁴⁰, R⁵⁴¹, R⁵⁴², R⁵⁴³, R⁵⁴⁴, R⁵⁴⁵, R⁵⁴⁶, R⁵⁴⁷, R⁵⁴⁸, R⁵⁴⁹, R⁵⁵⁰, R⁵⁵¹, R⁵⁵², R⁵⁵³, R⁵⁵⁴, R⁵⁵⁵, R⁵⁵⁶, R⁵⁵⁷, R⁵⁵⁸, R⁵⁵⁹, R⁵⁶⁰, R⁵⁶¹, R⁵⁶², R⁵⁶³, R⁵⁶⁴, R⁵⁶⁵, R⁵⁶⁶, R⁵⁶⁷, R⁵⁶⁸, R⁵⁶⁹, R⁵⁷⁰, R⁵⁷¹, R⁵⁷², R⁵⁷³, R⁵⁷⁴, R⁵⁷⁵, R⁵⁷⁶, R⁵⁷⁷, R⁵⁷⁸, R⁵⁷⁹, R⁵⁸⁰, R⁵⁸¹, R⁵⁸², R⁵⁸³, R⁵⁸⁴, R⁵⁸⁵, R⁵⁸⁶, R⁵⁸⁷, R⁵⁸⁸, R⁵⁸⁹, R⁵⁹⁰, R⁵⁹¹, R⁵⁹², R⁵⁹³, R⁵⁹⁴, R⁵⁹⁵, R⁵⁹⁶, R⁵⁹⁷, R⁵⁹⁸, R⁵⁹⁹, R⁶⁰⁰, R⁶⁰¹, R⁶⁰², R⁶⁰³, R⁶⁰⁴, R⁶⁰⁵, R⁶⁰⁶, R⁶⁰⁷, R⁶⁰⁸, R⁶⁰⁹, R⁶¹⁰, R⁶¹¹, R⁶¹², R⁶¹³, R⁶¹⁴, R⁶¹⁵, R⁶¹⁶, R⁶¹⁷, R⁶¹⁸, R⁶¹⁹, R⁶²⁰, R⁶²¹, R⁶²², R⁶²³, R⁶²⁴, R⁶²⁵, R⁶²⁶, R⁶²⁷, R⁶²⁸, R⁶²⁹, R⁶³⁰, R⁶³¹, R⁶³², R⁶³³, R⁶³⁴, R⁶³⁵, R⁶³⁶, R⁶³⁷, R⁶³⁸, R⁶³⁹, R⁶⁴⁰, R⁶⁴¹, R⁶⁴², R⁶⁴³, R⁶⁴⁴, R⁶⁴⁵, R⁶⁴⁶, R⁶⁴⁷, R⁶⁴⁸, R⁶⁴⁹, R⁶⁵⁰, R⁶⁵¹, R⁶⁵², R⁶⁵³, R⁶⁵⁴, R⁶⁵⁵, R⁶⁵⁶, R⁶⁵⁷, R⁶⁵⁸, R⁶⁵⁹, R⁶⁶⁰, R⁶⁶¹, R⁶⁶², R⁶⁶³, R⁶⁶⁴, R⁶⁶⁵, R⁶⁶⁶, R⁶⁶⁷, R⁶⁶⁸, R⁶⁶⁹, R⁶⁷⁰, R⁶⁷¹, R⁶⁷², R⁶⁷³, R⁶⁷⁴, R⁶⁷⁵, R⁶⁷⁶, R⁶⁷⁷, R⁶⁷⁸, R⁶⁷⁹, R⁶⁸⁰, R⁶⁸¹, R⁶⁸², R⁶⁸³, R⁶⁸⁴, R⁶⁸⁵, R⁶⁸⁶, R⁶⁸⁷, R⁶⁸⁸, R⁶⁸⁹, R⁶⁹⁰, R⁶⁹¹, R⁶⁹², R⁶⁹³, R⁶⁹⁴, R⁶⁹⁵, R⁶⁹⁶, R⁶⁹⁷, R⁶⁹⁸, R⁶⁹⁹, R⁷⁰⁰, R⁷⁰¹, R⁷⁰², R⁷⁰³, R⁷⁰⁴, R⁷⁰⁵, R⁷⁰⁶, R⁷⁰⁷, R⁷⁰⁸, R⁷⁰⁹, R⁷¹⁰, R⁷¹¹, R⁷¹², R⁷¹³, R⁷¹⁴, R⁷¹⁵, R⁷¹⁶, R⁷¹⁷, R⁷¹⁸, R⁷¹⁹, R⁷²⁰, R⁷²¹, R⁷²², R⁷²³, R⁷²⁴, R⁷²⁵, R⁷²⁶, R⁷²⁷, R⁷²⁸, R⁷²⁹, R⁷³⁰, R⁷³¹, R⁷³², R⁷³³, R⁷³⁴, R⁷³⁵, R⁷³⁶, R⁷³⁷, R⁷³⁸, R⁷³⁹, R⁷⁴⁰, R⁷⁴¹, R⁷⁴², R⁷⁴³, R⁷⁴⁴, R⁷⁴⁵, R⁷⁴⁶, R⁷⁴⁷, R⁷⁴⁸, R⁷⁴⁹, R⁷⁵⁰, R⁷⁵¹, R⁷⁵², R⁷⁵³, R⁷⁵⁴, R⁷⁵⁵, R⁷⁵⁶, R⁷⁵⁷, R⁷⁵⁸, R⁷⁵⁹, R⁷⁶⁰, R⁷⁶¹, R⁷⁶², R⁷⁶³, R⁷⁶⁴, R⁷⁶⁵, R⁷⁶⁶, R⁷⁶⁷, R⁷⁶⁸, R⁷⁶⁹, R⁷⁷⁰, R⁷⁷¹, R⁷⁷², R⁷⁷³, R⁷⁷⁴, R⁷⁷⁵, R⁷⁷⁶, R⁷⁷⁷, R⁷⁷⁸, R⁷⁷⁹, R⁷⁸⁰, R⁷⁸¹, R⁷⁸², R⁷⁸³, R⁷⁸⁴, R⁷⁸⁵, R⁷⁸⁶, R⁷⁸⁷, R⁷⁸⁸, R⁷⁸⁹, R⁷⁹⁰, R⁷⁹¹, R⁷⁹², R⁷⁹³, R⁷⁹⁴, R⁷⁹⁵, R⁷⁹⁶, R⁷⁹⁷, R⁷⁹⁸, R⁷⁹⁹, R⁸⁰⁰, R⁸⁰¹, R⁸⁰², R⁸⁰³, R⁸⁰⁴, R⁸⁰⁵, R⁸⁰⁶, R⁸⁰⁷, R⁸⁰⁸, R⁸⁰⁹, R⁸¹⁰, R⁸¹¹, R⁸¹², R⁸¹³, R⁸¹⁴, R⁸¹⁵, R⁸¹⁶, R⁸¹⁷, R⁸¹⁸, R⁸¹⁹, R⁸²⁰, R⁸²¹, R⁸²², R⁸²³, R⁸²⁴, R⁸²⁵, R⁸²⁶, R⁸²⁷, R⁸²⁸, R⁸²⁹, R⁸³⁰, R⁸³¹, R⁸³², R⁸³³, R⁸³⁴, R⁸³⁵, R⁸³⁶, R⁸³⁷, R⁸³⁸, R⁸³⁹, R⁸⁴⁰, R⁸⁴¹, R⁸⁴², R⁸⁴³, R⁸⁴⁴, R⁸⁴⁵, R⁸⁴⁶, R⁸⁴⁷, R⁸⁴⁸, R⁸⁴⁹, R⁸⁵⁰, R⁸⁵¹, R⁸⁵², R⁸⁵³, R⁸⁵⁴, R⁸⁵⁵, R⁸⁵⁶, R⁸⁵⁷, R⁸⁵⁸, R⁸⁵⁹, R⁸⁶⁰, R⁸⁶¹, R⁸⁶², R⁸⁶³, R⁸⁶⁴, R⁸⁶⁵, R⁸⁶⁶, R⁸⁶⁷, R⁸⁶⁸, R⁸⁶⁹, R⁸⁷⁰, R⁸⁷¹, R⁸⁷², R⁸⁷³, R⁸⁷⁴, R⁸⁷⁵, R⁸⁷⁶, R⁸⁷⁷, R⁸⁷⁸, R⁸⁷⁹, R⁸⁸⁰, R⁸⁸¹, R⁸⁸², R⁸⁸³, R⁸⁸⁴, R⁸⁸⁵, R⁸⁸⁶, R⁸⁸⁷, R⁸⁸⁸, R⁸⁸⁹, R⁸⁹⁰, R⁸⁹¹, R⁸⁹², R⁸⁹³, R⁸⁹⁴, R⁸⁹⁵, R⁸⁹⁶, R⁸⁹⁷, R⁸⁹⁸, R⁸⁹⁹, R⁹⁰⁰, R⁹⁰¹, R⁹⁰², R⁹⁰³, R⁹⁰⁴, R⁹⁰⁵, R⁹⁰⁶, R⁹⁰⁷, R⁹⁰⁸, R⁹⁰⁹, R⁹¹⁰, R⁹¹¹, R⁹¹², R⁹¹³, R⁹¹⁴, R⁹¹⁵, R⁹¹⁶, R⁹¹⁷, R⁹¹⁸, R⁹¹⁹, R⁹²⁰, R⁹²¹, R⁹²², R⁹²³, R⁹²⁴, R⁹²⁵, R⁹²⁶, R⁹²⁷, R⁹²⁸, R⁹²⁹, R⁹³⁰, R⁹³¹, R⁹³², R⁹³³, R⁹³⁴, R⁹³⁵, R⁹³⁶, R⁹³⁷, R⁹³⁸, R⁹³⁹, R⁹⁴⁰, R⁹⁴¹, R⁹⁴², R⁹⁴³, R⁹⁴⁴, R⁹⁴⁵, R⁹⁴⁶, R⁹⁴⁷, R⁹⁴⁸, R⁹⁴⁹, R⁹⁵⁰, R⁹⁵¹, R⁹⁵², R⁹⁵³, R⁹⁵⁴, R⁹⁵⁵, R⁹⁵⁶, R⁹⁵⁷, R⁹⁵⁸, R⁹⁵⁹, R⁹⁶⁰, R⁹⁶¹, R⁹⁶², R⁹⁶³, R⁹⁶⁴, R⁹⁶⁵, R⁹⁶⁶, R⁹⁶⁷, R⁹⁶⁸, R⁹⁶⁹, R⁹⁷⁰, R⁹⁷¹, R⁹⁷², R⁹⁷³, R⁹⁷⁴, R⁹⁷⁵, R⁹⁷⁶, R⁹⁷⁷, R⁹⁷⁸, R⁹⁷⁹, R⁹⁸⁰, R⁹⁸¹, R⁹⁸², R⁹⁸³, R⁹⁸⁴, R⁹⁸⁵, R⁹⁸⁶, R⁹⁸⁷, R⁹⁸⁸, R⁹⁸⁹, R⁹⁹⁰, R⁹⁹¹, R⁹⁹², R⁹⁹³, R⁹⁹⁴, R⁹⁹⁵, R⁹⁹⁶, R⁹⁹⁷, R⁹⁹⁸, R⁹⁹⁹, R¹⁰⁰⁰, R¹⁰⁰¹, R¹⁰⁰², R¹⁰⁰³, R¹⁰⁰⁴, R¹⁰⁰⁵, R¹⁰⁰⁶, R¹⁰⁰⁷, R¹⁰⁰⁸, R¹⁰⁰⁹, R¹⁰¹⁰, R¹⁰¹¹, R^{1012</}

FOR THE PURPOSES OF INFORMATION ONLY

Codes used to identify States party to the PCT on the front pages of pamphlets publishing international applications under the PCT.

AT Austria
AU Australia
BB Barbados
BE Belgium
BG Bulgaria
BT Bhutan

FR France
GA Gabon
GB United Kingdom
HU Hungary
IT Italy
JM Jamaica

ML Mali
MR Mauritania
MW Malawi
NL Netherlands
NO Norway
PR Puerto Rico

PT Portugal
RO Romania
RU Russian Federation
DE Germany, Federal Republic of
DK Denmark
FI Finland

LU Luxembourg
MC Monaco
MG Madagascar
MT Malta

IG Togo
US United States of America
VN Vietnam

- 1 -

PROTEASE INHIBITORS

Background of the Invention

This invention relates to protease inhibitors, and particularly to transition state analogs.

5 Transition state analogs, compounds which are thought to resemble the substrates of enzymes, are thought to bind more tightly to the enzymes than the substrates themselves. Transition state analogs form complexes with enzymes at their catalytic sites.


10 Baugh et al. (Proteinases and Tumor Invasion ed. Strauli et al., Raven Press, New York, 1980, p./165) state that transition state analogs containing boronic acid moieties or aldehydes form tetrahedral adducts with serine proteases and are thus good inhibitors of these
15 enzymes. Further, they state that some peptide aldehydes have been synthetically prepared, that most are of microbial origin, and that "it would appear that changing the R-group [of synthetic peptides] to satisfy the specific requirements of a given protease should
20 result in both potent and specific inhibitors." They also state that transition state analogs containing cyclic ester moieties have been used to inhibit chymotrypsin and that "variations thereof may become useful as inhibitors of cathepsin G."

25 Yoshimoto et al. (J. Biochem. 98: 975, 1985) describe prolyl endopeptidase inhibitors containing a protinal moiety. These inhibitors appear to act non-competitively.

Shenvi et al., U.S. Patent No. 4,499,082,
30 describe peptides having an α -amino boronic acid

- 2 -

residue. These peptides are reversible inhibitors of elastase. They have the structural formula



where R_2 is an alkyl group of one to six carbons which
5 may have an aromatic substituent or an in-chain bivalent group.


Summary of the Invention

In a first aspect, the invention features
compounds having the structure


10



and salts thereof, where T is a boronate group of the formula



where each D^1 and D^2 , independently, is a hydroxyl
15 group or a group which is capable of being hydrolyzed to a hydroxyl group in aqueous solution at physiological pH; or T is a group of the formula






where G is either H, F or an alkyl group containing 1 to
20 about 20 carbon atoms and optional heteroatoms which can be N, S, or O; or T is a phosphonate group of the formula



- 3 -

alkyl (each containing about 1-20 carbon atoms) and, optionally, heteroatoms which can be N, S, or O; where T is a group able to form a complex with the catalytic site of an enzyme); X includes one or more amino acids,
 5 Y is




and each R^1 , R^2 , R^3 , R^4 , R^5 , R^6 , R^7 , and
 10 R^8 , independently, is a group which does not interfere significantly (i.e., does not raise the K_i of the compound to greater than $10^{-5}M$) with site-specific recognition of the compound by the enzyme, while
 15 permitting a complex to be formed between the compound and the enzyme.

In preferred embodiments, T is a boronate group, each D^1 and D^2 , independently, is OH or F or D^1 and D^2 together form a ring containing 1 to about 20 carbon atoms, and optionally heteratoms which can be
 20 N, S, or O; each R^{1-8} is H; X mimics the substrate recognised by the enzyme, for example X is pro-, thr-pro-, ala-pro-, ala-ala-pro-, ser-thr-pro-, pro-ser-, pro-thr-, or ser-pro- (pro = proline, thr = threonine, and ser = serine); X contains both an amino
 25 acid and a blocking group, such as an acetyl group; the

- 4 -

proteases, even more preferably IgA1 proteases; and the analog has a binding constant of at least $10^{-7}M$, most preferably $10^{-10}M$.

In a second aspect, the invention features a compound, which is useful as an intermediate in the synthesis of compounds of Formula (1), having the formula



where V is $(\text{CH}_3)_3\text{Si}-$ or $\text{H}-$, Y is



10



or



each R^1 , R^2 , R^3 , R^4 , R^5 , R^6 , R^7 , and R^8 is independently H, or C_1 - C_{10} alkyl or aryl, and Z is



15 where D^1 and D^2 are as defined above.

The invention also features a method for producing the above compounds. The method includes the steps of reacting a compound of the formula

- 5 -

where Q is Cl or Br, and R², Y, Z, and D¹ and D² are as defined above, with a compound of the formula



in an inert solvent at a temperature between -78°C and 5 25°C, and heating the reaction mixture to at least 80°C.

In preferred embodiments the reacting step is performed in the absence of an inert solvent.

The compounds of the invention are peptide derivatives (or intermediates in their formation) which 10 are potent inhibitors of proteolytic enzymes (especially those produced by pathogens) which are post-prolyl cleaving enzymes, e.g., the serine proteases able to act on IgA1 proteins. The compounds generally have a proline, proline analog, a 2-azetidinecarboxylic acid, 15 or pipecolic acid linked to a group, for example a boronate group, which mimics the transition state of an enzyme substrate, and to a peptide moiety which preferably mimics the site of the substrate acted upon by the enzyme of interest. It is proposed that the 20 peptide moiety is recognized as a substrate by the enzyme to be inhibited, and it then enters an active site, catalytic site, or transition state binding site of the enzyme, and the transition state-mimicing group of the compound of the invention binds strongly at this 25 site. This binding advantageously prevents the enzyme from acting upon its natural substrate. The high affinity of these compounds make them effective at concentrations as low as 10⁻⁷M, or even 10⁻¹⁰M.

Thus, the present invention also provides 30 compositions including one or more compounds of formula

- 6 -

Other features and advantages of the invention will be apparent from the following description of preferred embodiments thereof, and from the claims.

Description of the Preferred Embodiments

5 The Figures will first briefly be described.

Drawing

Figure 1 shows the general formula of three preferred compounds.

10 Figure 2 is a diagrammatic representation of the synthesis of a boroProline compound.

Structure

The compounds of the invention have the general structure recited in the Summary of the Invention above. Examples of preferred structures are those
15 referred to as preferred embodiments above.

The structure of the compounds is such that at least a portion of the amino acid sequence near the cleavage site of a post-prolyl cleaving enzyme substrate is duplicated, or nearly duplicated. This duplication
20 is in part responsible for the ability of the compounds to inhibit the enzyme, by a mechanism thought to involve competitive inhibition between the compound and the actual enzyme substrate.


The choice of amino acid sequence affects the
25 inhibitory activity of the compound, and its specificity. As a first step in determining a suitable sequence, the amino acid sequence of the substrate is determined near its cleavage site. In some cases, such as for serine proteases, a suitable inhibitor sequence
30 is the amino acid sequence N-terminal to (i.e., to the left of) the proline cleavage point, and includes that proline. Peptide fragments can be synthesized and then

- 7 -


similar fashion, by testing the inhibitory effect of a particular compound on a variety of enzyme activities. The compounds preferably inhibit the activity of enzymes detrimental to an animal or human patient, and
 5 preferably do not inhibit necessary enzymes.

The compounds of the invention can be designed so that they are resistant to attack by agents which might otherwise cause their catabolic degradation by cleavage of one or more peptide bonds in the peptides.
 10 For example, an N-terminal blocking group, such as acetyl, can increase the half-life of the compounds in vivo, and thus improves inhibition.

The structure of such blocking groups can vary widely. In one blocking reaction, a hydrogen atom of
 15 the amino terminal amino group is replaced, generally in a dehydration reaction. Thus, blocking groups such as



readily added to a peptide chain. Others include



20 Such N-terminal blocking groups may be employed not only to protect amino-terminal groups, but also may protect-side chain amino group of amino acids which make up X, such as where X includes Lys or Arg. Similarly, amino acid residues having acidic or hydroxy side chains
 25 can be protected in the form of t-butyl, benzyl, or

increased half-life

- 8 -

The compounds also include a group (T) which causes the compound to complex with an enzyme, not only in a competitive fashion, but in a chemically reactive manner to form a strong bond between the compound and the enzyme. This group thus acts to bind the compound to the enzyme, and increases the inhibitory binding constant (K_i) of the compound. Examples of such groups include boronates, fluoroalkyl ketones and substituted phosphonates (of the formulae given in the Summary above, examples of which are shown in Fig. 1). These groups are covalently bonded to the prolyl residue of the compound, as in the above formula.

The proline or proline analog, represented by



above, is chosen so that it mimics the structure of proline recognized by the active site of the enzyme to be inhibited. It can be modified by providing R^{1-2} groups which do not interfere significantly with this recognition and thus do not significantly affect the K_i of a compound. Thus, one or more hydroxyl groups can be substituted to form hydroxy-proline, and methyl or sugar moieties may be linked to these groups. One skilled in the art will recognize that these groups are not critical in this invention and that a large choice of substituents are acceptable for R.

Examples of compounds having utility as serine protease inhibitors include compounds having a peptide chain similar to the substrate of IgA1 and IgA2

- 9 -

bonds of IGA, the immunoglobulin that provides antibody defense of mucosal surfaces, resulting in a nonfunctional immunoglobulin and impairment of the host defense system. This is expected to be a strong
5 contributing factor to pathogenesis of organisms such as Streptococcus sanguis, S. pneumoniae, Neisseria gonorrhoeae, N. meningitidis, and Haemophilus influenzae.

IgA1 proteases recognize the cleavage site Ser-Thr-Pro-Pro-X (where X is any amino acid),
10 hydrolyzing between Pro and X (i.e., they are post-prolyl cleaving enzymes). Accordingly, Ser-Thr-Pro-Pro-T is a suitable compound of the invention for inhibiting IGA1 proteases. The Ser or Thr in this compound can be readily substituted with any of
15 the 20 naturally occurring amino acids, most preferably those having non-bulky side groups, such as Ala and Gly. It is also possible to substitute non-naturally occurring amino acids, such as 2-azetidinecarboxylic acid or pipecolic acid (which have six-membered, and
20 four-membered ring structures respectively) for either of the Pro residues. Those skilled in the art will recognize that there are other such changes which can be made without significantly affecting the inhibitory character of these compounds.

25 In the case of type 1 IgA proteases, the cleavage site is Pro-Thr-Pro-X, with hydrolysis occurring between Pro and X. Thus, a preferred compound of the invention for inhibiting IGA2 proteases has the formula Pro-Thr-Pro-T. Thr can be substituted by any of
30 the naturally occurring amino acids, especially ones having non-bulky side groups, such as Ala, Gly, or Ser.

Other examples of enzymes which can be

- 10 -

degrading enzymes, vasopressin degrading enzymes, and oxytocin degrading enzymes. Further, the serine protease, dipeptidyl peptidase Type IV, on T-lymphocytes (Andrews et al., Clin. Lab. Haemato. 7: 359-368, 1985),
5 which plays a role in regulation of the immune response, can be inhibited by suitable such compounds. These inhibitors may be useful in treatment of AIDS. Walters et al., Molecular and Cellular Biochemistry 30: 111-127, 1980 describes other such enzymes, and is hereby
10 incorporated by reference.

Synthesis

Synthesis of boroProline

Referring to Fig. 2, the starting compound I is prepared essentially by the procedure of Matteson et
15 al., 3 Organometallics 1284, 1984, except that a pinacol ester is substituted for the pinanediol ester. Similar compounds such as boropipericolic acid and 2-azetidine boronic acid can be prepared by making the appropriate selection of starting material to yield the pentyl and
20 propyl analogs of compound I. Further, Cl can be substituted for Br in the formula, and other diol protecting groups can be substituted for pinacol in the formula, e.g., 2,3-butanediol and alpha-pinanediol.

Compound II is prepared by reacting compound I
25 with $[(CH_3)_3Si]_2N^-Li^+$. In this reaction hexamethyldisilazane is dissolved in tetrahydrofuran and an equivalent of n-butyllithium added at $-78^\circ C$. After warming to room temperature ($20^\circ C$) and cooling to $-78^\circ C$ an equivalent of compound I is added in
30 tetrahydrofuran. The mixture is allowed to slowly come to room temperature and to stir overnight. The alpha-bis[trimethylsilane]-protected amine is isolated

- 11 -

under a nitrogen blanket, yielding a hexane solution of compound 2.

Compound III, the N-trimethysilyl protected form of boroProline is obtained by the thermal
5 cyclization of compound II during the distillation process in which compound II is heated to 100-150°C and distillate is collected which boils 66-62°C at 0.06-0.10 mm pressure.

Compound IV, boroProline-pinacol hydrogen
10 chloride, is obtained by treatment of compound III with HCl:dioxane. Excess HCl and by-products are removed by trituration with ether. The final product is obtained in a high degree of purity by recrystallization from ethyl acetate. H-boroProline as the hydrochloride salt
15 is preferred, but other salts forms such as the hydrobromide and trifluoroacetate can readily be obtained by substitution of the appropriate acid for HCl.

The boroProline esters can also be obtained by treatment of the reaction mixture obtained in the
20 preparation of compound II with anhydrous acid to yield 1-amino-4-bromobutyl boronate pinacol as a salt. Cyclization occurs after neutralizing the salt with base and heating the reaction.

Synthesis of boroProline Peptides

25 General methods of coupling of N-protected peptides and amino acids with suitable side-chain protecting groups to H-boroProline-pinacol are applicable. When needed, side-chain protecting and N-terminal protecting groups can be removed by treatment
30 with anhydrous HCl, HBr, trifluoroacetic acid, or by catalytic hydrogenation. These procedures are known to those skilled in the art.

- 12 -

protecting groups from threonine hydroxyl group results in a complex mixture of products. Thus, the use of hydrogenolytic protecting groups for threonyl residue is preferred.

5 The mixed anhydride procedure of Anderson et al., J. Am. Chem. Soc., 89:5012 (1984) is preferred for peptide coupling. The mixed anhydride of an N-protected amino acid or a peptide varying in length from a dipeptide to tetrapeptide is prepared by dissolving the
10 peptide in tetrahydrofuran and adding one equivalent of N-methylmorpholine. The solution is cooled to -20°C and an equivalent of isobutyl chloroformate is added. After 5 minutes, this mixture and one equivalent of triethylamine (or other sterically hindered base) are
15 added to a solution of H-boroPro-pinacol dissolved in either cold chloroform or tetrahydrofuran.

 The reaction mixture is routinely stirred for one hour at -20°C and 1-2 hours at room temperature (20°C). Solvent is removed by evaporation, and the
20 residue is dissolved in ethyl acetate. The organic solution is washed with 0.20 N hydrochloric acid, 5% aqueous sodium bicarbonate, and saturated aqueous sodium chloride. The organic phase is dried over anhydrous sodium sulfate, filtered, and evaporated. Products are
25 purified by either silica gel chromatography or gel permeation chromatography using Sephadex™ LH-20 and methanol as a solvent.

 Previous studies have shown that the pinacol protecting group can be removed in situ by preincubation
30 in phosphate buffer prior to running biological experiments; Kettner et al., J. Biol. Chem. 259: 15100-15114 (1984). Several other methods are also

- 13 -

product. First, the peptide can be treated with diethanolamine to yield the corresponding diethanolamine boronic acid ester, which can be readily hydrolyzed by treatment with aqueous acid or a sulfonic acid substituted polystyrene resin as described in Kettner et al., Id. Both pinacol and pinanediol protecting groups can be removed by treating with BCl_3 in methylene chloride as described by Kinder et al., J. Med. Chem., 28: 1917. Finally, the free boronic acid can be converted to the difluoroboron derivative ($-\text{BF}_2$) by treatment with aqueous HF as described by Kinder et al., Id.

Similarly, different ester groups can be introduced by reacting the free boronic acid with various di-hydroxy compounds (for example, those containing heteroatoms such as S or N) in an inert solvent.

The following abbreviations are used in the examples below. THF - tetrahydrofuran; H-Pro-OBzl - the benzyl ester of proline; H-Thr (OBzl)-OH - the benzyl ether derivative of threonine; Boc - the tertiary butyloxycarbonyl group; FABMS - fast atom bombardment mass spectrometry.

All natural amino acids are in the L-configuration. H-boroProline is in the D,L-configuration.

Example 1: Preparation of boroProline-pinacol
(H-boroPro-pinacol)

The intermediate, 4-Bromo-1-chlorobutyl boronate pinacol, was prepared by the method in Matteson et al., Organometallics, (3): 1284-1288 (1984), except

pinacol protecting group

- 14 -

3-bromopropyl boronate pinacol was prepared by hydrogenboration of allyl bromide (173 ml, 2.00 moles) with catechol borane (240 ml, 2.00 moles). Catechol borane was added to allyl bromide and the reaction
5 heated for 4 hours at 100°C under a nitrogen atmosphere. The product, 3-bromopropyl boronate catechol (bp 95-102°C, 0.25 mm), was isolated in a yield of 49% by distillation. The catechol ester (124 g, 0.52 moles) was transesterified with pinacol (61.5 g, 0.52
10 moles) by mixing the component in 50 ml of THF and allowing them to stir for 0.5 hours at 0°C and 0.5 hours at room temperature. Solvent was removed by evaporation and 250 ml of hexane added. Catechol was removed as a crystalline solid. Quantitative removal was achieved by
15 successive dilution to 500 ml and to 1000 ml with hexane and removing crystals at each dilution. Hexane was evaporated and the product distilled to yield 177 g (bp 60-64°C, 0.35 mm).

4-Bromo-1-chlorobutyl boronate pinacol was
20 prepared by homologation of the corresponding propyl boronate. Methylene chloride (50.54 ml, 0.713 moles) was dissolved in 500 ml of THF, 1.54 N n-butyllithium in hexane (480 ml, 0.780 moles) was slowly added at -100°C. 3-Bromopropyl boronate pinacol (178 g, 0.713
25 moles) was dissolved in 500 ml of THF, cooled to the freezing point of the solution, and added to the reaction mixture. Zinc chloride (54.4 g, 0.392 moles) was dissolved in 250 ml of THF, cooled to 0°C, and added to the reaction mixture in several portions. The
30 reaction was allowed to slowly warm to room temperature and to stir overnight. Solvent was evaporated and the residue dissolved in hexane (1 liter) and washed with

- 15 -

filtering, solvent was evaporated. The product was distilled to yield 147 g (bp 110-112°C, 0.200 mm).

N-Trimethylsilyl-boroProline pinacol was prepared first by dissolving hexamethyldisilazane (20.0 g, 80.0 mmoles) in 30 ml of THF, cooling the solution to -78°C, and adding 1.62 N n-butyllithium in hexane (49.4 ml, 80.0 mmoles). The solution was allowed to slowly warm to room temperature. It was recooled to -78°C and 4-bromo-1-chlorobutyl boronate pinacol (23.9 g, 80.0 mmoles) added in 20 ml of THF. The mixture was allowed to slowly warm to room temperature and to stir overnight. Solvent was removed by evaporation and dry hexane (400 ml) added to yield a precipitant which was removed by filtration under an nitrogen atmosphere. The filtrate was evaporated and the residue distilled, yielding 19.4 g of the desired product (bp 60-62°C, 0.1-0.06 mm).

H-boroProline-pinacol.HCl was prepared by cooling N-trimethylsilyl-boroProline-pinacol (16.0 g, 61.7 mmoles) to -78°C and adding 4 N HCL:dioxane 46 ml, 185 mmoles). The mixture was stirred 30 minutes at -78°C and 1 hour at room temperature. Solvent was evaporated and the residue triturated with ether to yield a solid. The crude product was dissolved in chloroform and insoluble material removed by filtration. The solution was evaporated and the product crystallized from ethyl acetate to yield 11.1 g of the desired product (mp 156.5-157°C).

Example 2: Preparation of Boc-Ala-Pro-boroPro-pinacol

Boc-Ala-Pro-boroPro-pinacol was prepared by coupling Boc-Ala-Pro-OH to H-boroPro-pinacol. First,

- 16 -

52.8 mmole) and isobutyl chloroformate (6.8 ml, 52.8 mmole) for 5 minutes in 50 ml of THF at -20° . The reaction mixture and additional N-methylmorpholine (5.8 ml) were added to H-Pro-OBzl.HCl (12.8 g, 52.8 mmole) dissolved in 50 ml of cold chloroform. After the mixture was stirred for 1 hour at -20°C and 2 hours at room temperature, it was filtered and the filtrate evaporated. The residue was dissolved in ethyl acetate and washed sequentially with 0.2 N hydrochloric acid, 5% aqueous NaCO_3 , and saturated aqueous NaCl. The organic phase was dried over anhydrous Na_2SO_4 , filtered, and evaporated to yield Boc-Ala-Pro-OBzl as an oil (14.1 g). The benzyl ester protecting group was removed by dissolving Boc-Ala-Pro-OBzl (35 g, 92.8 mmole) in 200 ml of methanol and hydrogenating for 1 hour in the presence of 0.5 g of 10% Pd/C. The catalyst was removed by filtration and the solvent evaporated. The residue was crystallized from ethyl acetate to yield 16.1 g of Boc-Ala-Pro-OH (mp $153.5\text{--}154.5^{\circ}\text{C}$).

Boc-Ala-Pro-OH (1.26 g, 4.28 mmole) was coupled to H-boroPro-pinacol by the general procedure described for the preparation of Boc-Ala-Pro-OBzl. Boc-Ala-Pro-OH (1.26 g, 4.28 mmole) was dissolved in 10 ml of THF and cooled to -20°C ; N-methylmorpholine (0.47 ml, 4.28 mmole) and isobutyl chloroformate (0.557 ml, 4.28 mmole) were added. After stirring for 5 minutes, 10 ml of cold THF and triethylamine (0.597 ml, 4.28 mmole) were added and the mixture added to a cold solution of H-boroPro-pinacol.HCl (1.0 g, 4.28 mmole) in 5 ml chloroform. After dissolving the reaction product in ethyl acetate and washing with aqueous HCl, NaHCO_3 , and saturated aqueous NaCl, 0.39 g of material were

- 17 -

Example 3: Preparation of H-Ala-Pro-
boroPro-pinacol.HCL

Boc-Ala-Pro-boroPro-pinacol (0.58 g, 1.25
mmoles) was allowed to react with 2.5 ml of 4 N
5 HCl:dioxane for 30 minutes at room temperature. Ether
(50 ml) was added to yield 0.22 g of amorphous white
solid.

Example 4: Preparation of Ac-Ala-Pro-boroPro-pinacol

H-Ala-Pro-boroPro-pinacol.HCL (0.22 g, 0.55
10 mmoles) was dissolved in 3 ml of THF and cooled to 0°C.
Acetic anhydride (0.078 ml, 0.825 mmoles) and
triethylamine (0.115 ml, 0.825 mmoles) were added and
the reaction was allowed to come to room temperature.
After approximately 25 minutes, additional triethylamine
15 (0.25 ml, 0.179 mmoles) was added. After a total
reaction time of 45 minutes, the reaction solution was
applied to a 2.5 X 100 column of LH-20 in methanol and
fractions (approximately 7 ml) collected. Fraction
22-28 contained 0.16 g of the desired product.

20 Example 5: Preparation of MeOSuc-Ala-
Ala-Pro-boroPro-pinacol.

MeOSuc-Ala-Ala-Pro-OH was prepared by the
procedure described in Kettner et al., J. Biol. Chem.,
259: 15106-15114 (1984). MeOSuc-Ala-Ala-Pro-OH (1.59 g,
25 4.28 mmoles) was coupled to H-boroPro-pinacol.HCL (1.00
g, 4.28 mmoles) by the mixed anhydride procedure
described for the preparation of
Boc-Ala-Pro-boroPro-pinacol except that, after
filtration and evaporation of the reaction solvent, it
30 was applied to a 2 cm column containing 10 g of silica
gel equilibrated with chloroform. The column was eluted
with chloroform and fractions containing the desired

- 18 -

Example 6: Preparation of Boc-Pro-
Thr(OBzl)-boroPro-pinacol.

Boc-Pro-Thr(OBzl)-OH was prepared by coupling Boc-Pro-OSu (the N-hydroxysuccinamide ester of
5 Boc-Pro-OH) to H-Thr (OBzl) - OH, H-Thr (OBzl)-OH.HCL was dissolved in 25 ml of water, NaHCO_3 (6.21 g, 73.9 mmoles) and a solution of Boc-Pro-OSu (5.07 g, 16.25 mmoles) in 25 ml of dioxane were added. After stirring 3 hours, the reaction mixture was acidified with
10 hydrochloric acid and the product extracted into ethyl acetate. It was washed with 0.2 N HCl prepared in saturated aqueous NaCl, saturated NaCl and dried over anhydrous Na_2SO_4 . Solvent was evaporated to yield 6.24 g (mp 120-12.5°).

15 Boc-Pro-Thr(OBzl)-boroPro-pinacol was prepared by coupling Boc-Pro-Thr(OBzl)-OH (2.70 g, 6.42 mmoles) to H-boroPro-pinacol.HCL(1.50 g, 6.42 mmoles) using the procedure described for Boc-Ala-Pro-boroPro-pinacol. The product (2.4 g) was purified by chromatography on a
20 2.5 X 50 cm column of LH-20 in methanol and was obtained as 0.84 g of oil.

Example 7: Preparation of Boc-Pro-Thr-boroPro-Pinacol

Boc-Pro-Thr-boroPro-pinacol was prepared by hydrogenation of Boc-Pro-Thr(OBzl)-boroPro-pinacol (from
25 Example 6, 0.585 g, 0.79 mmoles). The protected peptide was dissolved in 100 ml of methanol and was hydrogenated for 2 hours on a Parr apparatus in the presence of 0.5 g of 10% Pd/C. The reaction was filtered and solvent evaporated. The product was allowed to stir with
30 hexane, and hexane insoluble material removed by filtration. The filtrate was evaporated to yield the desired product (0.24 g) as a white foam.

- 19 -

administered in an effective amount either alone or in combination with a pharmaceutically acceptable carrier or diluent.

The compounds of formula 1 or compositions
5 thereof can be used to treat mammals, e.g., humans, suffering from or subject to infections by pathogenic bacteria which produce deleterious post prolyl cleaving enzymes, e.g., the IgA1 protease of Haemophilus influenza. The compounds or compositions can be
10 administered alone or in combination with one another, or in combination with other therapeutic agents. The dosage level may be 1-500 mg/kg/day of the formula 1 compounds.


When administered to mammals (e.g., orally,
15 topically, intramuscularly, intraperitoneally, intravenously, parenterally, nasally or by suppository), the compounds of the invention enhance the ability of e.g., the immune system of the mammal to fight bacteria which produce an IgA-1 protease, thus slowing the course
20 of infection by the bacteria.

Other embodiments are within the following claims.

- 20 -

Claims

1. A compound, having the structure




wherein T is a group of the formula



where each D^1 and D^2 , independently, is a hydroxyl group or a group which is capable of being hydrolysed to a hydroxyl group in aqueous solution at physiological pH; a group of the formula



where G is either H, F or an alkyl group containing 1 to about 20 carbon atoms and optional heteroatoms which can be N, S, or O; or a phosphonate group of the formula






- 15 where each J, independently, is O-alkyl, N-alkyl, or alkyl, each said O-alkyl, N-alkyl or alkyl comprising about 1-20 carbon atoms and, optionally, heteroatoms which can be N, S, or O; said T being able to form a complex with the catalytic site of a post-prolyl
20 cleaving enzyme;

X comprises an amino acid,

- 21 -

Y is



- 5 and each R^1 , R^2 , R^3 , R^4 , R^5 , R^6 , R^7 , and R^8 , separately is a group which does not significantly interfere with site specific recognition of said compound by said enzyme, and allows said complex to be formed with said enzyme.
- 10 2. The compound of claim 1, wherein T is a boronate group.
3. The compound of claim 1, wherein T is a substituted phosphonate group or a fluoroalkyl ketone group.
- 15 4. The compound of claim 1 wherein each R^{1-8} is H.
5. The compound of claim 1 wherein X mimics the site of a substrate recognized by said enzyme.
6. The compound of claim 1 wherein X comprises
- 20 a blocking group.
7. The compound of claim 6 wherein X comprises an acetyl group.
8. The compound of claim 5 where X is pro-, thr-pro-, ala-pro-, ala-ala-pro-, ser-thr-pro-,
- 25 pro-ser-, pro-thr-, or ser-pro-.

- 22 -

10. The compound of claim 9, said enzyme being a serine protease.


11. The compound of claim 8, said enzyme being an IgA1 protease.

5 12. The compound of claim 1, wherein said compound has a binding constant to said enzyme of at least $10^{-10}M$.



13. The compound of claim 10, wherein said compound has a binding constant of at least $10^{-7}M$.

10 14. The compound of claim 1 admixed within a pharmaceutically acceptable carrier substance.

15. A compound having the formula



where V is $(\text{CH}_3)_3\text{Si}-$ or $\text{H}-$, Y is




each R^1 , R^2 , R^3 , R^4 , R^5 , R^6 , R^7 , and R^8 is independently H, or C_1 - C_{20} alkyl or aryl, and Z is




- 23 -

16. The compound of claim 1 or 14 wherein, each D^1 and D^2 is, independently, F or D^1 and D^2 together are a ring containing 1 to about 20 carbon atoms, and optionally heteroatoms which can be N, S, 5 or O.

17. A method for producing the compound of claim 15, comprising the steps of reacting



where Q is Cl or Br, and R^2 is H or C_1-C_{10} alkyl 10 or aryl, with



in an inert solvent at a temperature between $-78^\circ C$ and $25^\circ C$, and

heating the reaction mixture to at least $80^\circ C$.


15 18. The method of claim 17 wherein said reacting is in the presence of an inert solvent.

19. A method for inhibiting a post prolyl cleaving enzyme in a mammal, comprising administering to said mammal an effective amount of a compound of claim 1.


20 20. The method of claim 19 wherein said amount is 1-500 mg/kg/day.

Abstract of the Disclosure


A compound having the structure



where T is of the formula $-\text{B}-\text{D}^1$, where each D^1 and
 5 D^2 , independently, is a hydroxyl group or a group
 which is capable of being hydrolysed to a hydroxyl group
 in aqueous solution at physiological pH; a group of the
 formula




10 where G is either H, F or an alkyl group containing 1 to
 about 20 carbon atoms and optional heteroatoms which can
 be N, S, or O; or a phosphonate group of the formula




where J is O-alkyl, N-alkyl, or alkyl, each comprising
 15 about 1-20 carbon atoms and, optionally, heteroatoms
 which can be N, S, or O; T being able to form a complex
 with the catalytic site of an enzyme, X is a group
 having at least one amino acid,


Y is



or




or




5 and each R^1 , R^2 , R^3 , R^4 , R^5 , R^6 , R^7 , and
 R^8 is separately a group which does not interfere
 significantly (i.e., does not lower the K_i of the
 compound to less than $10^{-7}M$) with site-specific
 recognition of the compound by the enzyme, and allows a
 10 complex to be formed with the enzyme.


1 / 2



Prolyl Boronate




Prolyl Trifluoro alkyl ketone




Prolyl phosphonate

FIG 1

2 / 2



4-bromo-1-chlorobutyl boronate pinacol



4-bromo-1[(bistrimethylsilyl) amino] butyl boronate pinacol



1-trimethylsilyl-boroProline pinacol
(III)IV
boroProline-pinacol-HCl

FIG 2

INTERNATIONAL SEARCH REPORT

International Application No. **PCT/US88/03506**

I. CLASSIFICATION OF SUBJECT MATTER (If several classification symbols apply, indicate all) *

According to International Patent Classification (IPC) or to both National Classification and IPC

INT. CL. 4th Ed. A61K 37/02; C07K 5/06, 5/08, 5/10; see attn:
US. CL. 514/18,19; 530/329,330,331; 548/517,518,533,540

II. FIELDS SEARCHED

Minimum Documentation Searched *

Classification System

Classification Symbols

US 514/18,19; 530/329,330,331; 548/517,518,533,540

Documentation Searched other than Minimum Documentation
to the Extent that such Documents are Included in the Fields Searched *

III. DOCUMENTS CONSIDERED TO BE RELEVANT ^{1,2}

Category *	Citation of Document, ^{1,2} with indication, where appropriate, of the relevant passages ^{1,2}	Relevant to Claim No. ^{1,2}
A	US, A, 4,652,552 (Kettner) Published 24 March 1987	1-20
A	US, A, 4,636,492 (Kettner) Published 13 January 1987	1-20
A	US, A, 4,644,055 (Kettner) Published 17 February 1987	1-20
A	US, A, 4,582,821 (Kettner) Published 15 April 1986	1-20
A	US, A, 4,499,082 (Shenvi) Published 12 February 1985	1-20

* Special categories of cited documents: ^{1,2}

"A" document defining the general state of the art which is not considered to be of particular relevance

"E" earlier document but published on or after the international filing date

"L" document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified)

"O" document referring to an oral disclosure, use, exhibition or other means

"P" document published prior to the international filing date but later than the priority date claimed

"T" later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention

"X" document of particular relevance: the claimed invention cannot be considered novel or cannot be considered to involve an inventive step

"Y" document of particular relevance: the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art.

"Δ" document member of the same patent family

IV. CERTIFICATION

International Searching Authority

ISA/US

Signature of Authorized Officer

DELBERT H. PHILLIPS